

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/94502/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Butler, Christopher C., Sterne, Jonathan A. C., Lawton, Michael, O'Brien, Kathryn, Wootton, Mandy, Hood, Kerenza ORCID: <https://orcid.org/0000-0002-5268-8631>, Hollingworth, William, Little, Paul, Delaney, Brendan C., Van Der Voort, Judith, Dudley, Jan, Birnie, Kate, Pickles, Timothy ORCID: <https://orcid.org/0000-0001-7743-0234>, Waldron, Cherry-Ann ORCID: <https://orcid.org/0000-0001-8465-2492>, Downing, Harriet, Thomas-Jones, Emma, Lises, Catherine, Rumsby, Kate, Durbaba, Stevo, Whiting, Penny, Harman, Kim, Howe, Robin, MacGowan, Alasdair, Fletcher, Margaret and Hay, Alastair D. 2016. Nappy pad urine samples for investigation and treatment of UTI in young children: the 'DUTY' prospective diagnostic cohort study. British Journal of General Practice 66 (648) , e516-e524. 10.3399/bjgp16X685873 file

Publishers page: <http://dx.doi.org/10.3399/bjgp16X685873>  
<<http://dx.doi.org/10.3399/bjgp16X685873>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**'Nappy pad' urine samples for guiding investigation and treatment of urinary tract infection (UTI) in young children: Findings from the 'DUTY' prospective diagnostic cohort study**

Professor Christopher C. Butler, FRCGP, Professor of Primary Care (1)  
Professor Jonathan AC Sterne, PhD, Professor of Medical Statistics and Epidemiology (2)  
Mr Michael Lawton, MPhil, Research Associate (2)  
Dr Kathryn O'Brien, PhD, Clinical Lecturer (3)  
Dr Mandy Wootton, Lead Scientist (4)  
Professor Kerenza Hood, PhD, Director (5)  
Prof William Hollingworth, PhD, Professor of Health Economics (2)  
Professor Paul Little, FRCGP, Professor of Primary Care Research (6)  
Professor Brendan C Delaney, MD, Chair in Medical Informatics and Decision Making (7)  
Dr Judith van der Voort, FRCPCH, Consultant Paediatrician (8)  
Dr Jan Dudley, PhD, Consultant Paediatric Nephrologist (9)  
Dr Kate Birnie, PhD, Research Fellow (2)  
Mr Timothy Pickles, MSc, Statistician (5)  
Dr Cherry-Ann Waldron, PhD, Trial Manager (5)  
Mrs Harriet Downing, MPhil, Trial Manager (10)  
Dr Emma Thomas-Jones, PhD, Research Fellow (5)  
Mrs Catherine Lises, MSc, Data Manager (5)  
Ms Kate Rumsby, MSc, Trial Manager (6)  
Mr Stevo Durbaba, MSc, e-Resources Developer (11)  
Dr Penny Whiting, PhD, Senior Research Fellow (12)  
Dr Kim Harman, Dhealth, Trial Manager (6)  
Dr Robin Howe, FRCPath, Lead Scientist (4)  
Professor Alasdair MacGowan, MD, Professor of Clinical Microbiology and Antimicrobial Therapeutics (13)  
Professor Margaret Fletcher, PhD (14,15)  
Professor Alastair D Hay, FRCGP, Professor of Primary Care (10)

**Affiliations:**

1. Nuffield Department of Primary Care Health Sciences, University of Oxford, New Radcliffe House, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6NW and GP, Cwm Taf University Health Board.
2. School of Social and Community Medicine, University of Bristol, Canynge Hall, 39 Whatley Road, Clifton, Bristol BS8 2PS
3. Division of Population Medicine, School of Medicine, Cardiff University, Neuadd Meirionnydd, Heath Park, Cardiff, CF14 4YS 2.
4. Specialist Antimicrobial Chemotherapy Unit, Public Health Wales Microbiology Cardiff, University Hospital Wales, Heath Park, Cardiff CF14 4XW
5. South East Wales Trials Unit (SEWTU), Centre for Trials Research, Cardiff University, 7th Floor Neuadd Meirionnydd, Heath Park, Cardiff, CF14 4YS
6. Primary Care and Population Sciences Division, University of Southampton, Southampton, SO15 5ST
7. Department of Surgery and Cancer, Imperial College, St Mary's Hospital, London W2 2NY
8. Department of Paediatrics and Child Health, University Hospital of Wales, Heath Park, Cardiff CF14 4XW
9. Bristol Royal Hospital for Children, University Hospitals Bristol, NHS Foundation Trust, Bristol, BS2 8BJ.
10. Centre for Academic Primary Care, NIHR School of Primary Care Research, School of Social and Community Medicine, University of Bristol, Canynge Hall, 39 Whatley Road, Clifton Bristol, BS8 2PS, UK
11. King's College London, Division of Health and Social Care Research, Department of Primary Care and Public Health Sciences, 7th Floor, Capital House, 42 Weston Street, London, SE1 3QD, UK
12. NIHR CLAHRC West, University Hospitals Bristol NHS Foundation Trust, 9th Floor, Whitefriars, Lewins Mead, Bristol BS1 2NTHouse, 42 Weston Street, London, SE1 3QD
13. North Bristol NHS Trust, Southmead Hospital, Westbury-on-Trym, Bristol BS10 5NB

*Nappy pad urine sampling 03 February 2016*

14. University of the West of England, Bristol, UK, BS16 1DD

15. University Hospitals Bristol, NHS Foundation Trust, Bristol, BS2 8AE

*Corresponding author:* Professor Christopher C Butler, Nuffield Department of Primary Care Health Sciences, University of Oxford, New Radcliffe House, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6NW, UK. Email [christopher.butler@phc.ox.ac.uk](mailto:christopher.butler@phc.ox.ac.uk)

## **Abstract**

### **Background**

Although sampling urine using nappy pads is preferred by parents and recommended when a clean catch sample can't be obtained, we do not know the added diagnostic utility of 'nappy pad' urine samples, nor the proportion that are contaminated.

### **Setting**

Acutely unwell children <5 years presenting to 233 primary care sites in England and Wales.

### **Method**

Logistic regression to identify independent associations of symptoms, signs and urine dipstick test results with UTI; diagnostic utility quantified as area under the receiver operator curves (AUROC). Nappy pad rule characteristics, AUROC, and contamination compared to findings from clean catch samples.

### **Results**

Nappy pad samples were obtained from 3205 children (82% <2 years; 48% female), and culture results available for 2277 (71.0%). 30 (1.3%) met our laboratory definition of UTI. Female gender, smelly urine, darker urine, and absence of nappy rash were independently associated with UTI, with an internally validated, coefficient model AUROC of 0.81 (0.87 for clean catch) that increased to 0.87 (0.90 for clean catch) with the addition of dipstick results. GPs' 'working diagnosis' had an AUROC 0.63 (95% CI 0.53 to 0.72). 12.2% of nappy pad and 1.8% of clean-catch samples were 'frankly contaminated' (risk ratio 6.66; 95% CI 4.95 to 8.96;  $p < 0.001$ ).

### **Conclusion**

Nappy pad urine culture results, with features that can be reported by parents and dipstick tests, can be clinically useful, but are less accurate and more often contaminated compared to clean catch urine culture results, which should be prioritised. Dipstick testing adds diagnostic accuracy.

### **Key words**

Urinary Tract Infections; Paediatrics; Diagnosis; Anti-Bacterial Agents;

**How this fits in**

Up to 80% of urinary tract infections (UTI) in young children presenting to primary care are missed. Timely and accurate diagnosis is essential to avoid over- and under- treatment and investigation. This is especially difficult in pre-verbal children who are not toilet trained, and present with undifferentiated symptoms. General practitioners use, and parents prefer, nappy pads for collecting urine from children who are still in nappies, but the clinical utility of data derived from nappy pad samples, the added value of dipstick testing, and the proportion of contaminated samples is not known. We found that culture results from urine obtained using nappy pads, together with features that can be reported by parents, can be clinically useful in identifying acutely unwell pre-school children presenting to primary care who have a UTI, but with less accuracy compared to clean catch sampling. However, contamination rates are nearly seven times higher in nappy pad than in clean catch samples. Clean catch urine sampling in children in primary care should therefore be prioritised over the nappy pad method, but if urine sampling is done using nappy pads, then the addition of dipstick testing significantly improves diagnostic accuracy.

## **Introduction**

Urinary tract infection (UTI) may be missed in up to 80% of children presenting to primary care.(1, 2) Accurate diagnosis of UTI is essential to avoid over or under treatment with antibiotics and to appropriately target burdensome and expensive investigations.(3) This is especially important in younger, pre-verbal children who are not yet toilet trained and who often present with non-specific symptoms, making the decision about which children to investigate for UTI difficult.(3) Obtaining a urine sample can be time consuming and especially challenging in primary care where most children first present.(4) The nappy pad sampling method has been recommended by National Institute for Health and Care Excellence (NICE) in young children in nappies (diapers) when a clean catch sample cannot be obtained.(3) Urine sampling needs to be simple, reliable and acceptable, and parents find nappy pads easy to use, comfortable for their children, and prefer them to the clean catch method.(5) Nappy pad sampling is used in everyday care(1), and general practitioners report using nappy pad urine collection in over 40% of infants.(6) Many parents feel that the clean catch method is messy and time consuming and give up trying.(3, 5) However, the clinical utility of the information obtained from the nappy pad method of urine sampling is unclear, contamination rates may be higher than other sampling methods, and children in nappies present differently to older children who are more able to describe symptoms and in whom clean catch sampling is easier. Obtaining urine samples by more invasive methods such as supra-pubic aspiration or catheterisation is neither feasible nor acceptable in most primary care settings. We therefore aimed to develop a clinical prediction rule for the diagnosis of UTI based on sampling using the nappy pad method, and compare its diagnostic utility to a similar rule based on 'clean catch' urine samples (to be reported elsewhere in full).(7) In addition, we estimated the added diagnostic value of dipstick testing once a nappy pad sample was obtained and compared contamination rates by sampling method.

## **Methods**

### *Participants*

The Diagnosis of Urinary Tract infection in Young children (DUTY) study was a multicentre, prospective, diagnostic cohort study that recruited children aged less than five years in primary care.(8) Children were eligible if presenting with any acute (<28 days), undifferentiated illness (even when the clinician was confident of the diagnosis, such as a child with bronchiolitis) and/or new urinary symptoms.

### *Index tests and urine collection*

Following consent, 107 index test items were recorded. Parent reported items included the child's medical history and symptoms. Clinician assessed items came from a full clinical examination, including their global impression of illness severity (rated 0 to 10), their rating of the likelihood of UTI, and urine dipstick results

(performed after rating UTI likelihood). Index test items were derived from literature review and input from our co-investigator group.

We used the NICE recommended 'Newcastle Nappy Pads' for those children who wore nappies (diapers) and for those in whom the parent/guardian did not think clean catch would be successful.(3, 8, 9) First, the parent was asked to clean the nappy area using water or wipes (the wipes being supplied by the study). A nappy pad was inserted inside a clean nappy, and the nappy refastened. The nappy pad was removed as soon as the child urinated in order to reduce the risk of contamination. The perineum was cleaned again and a fresh pad inserted every 30 minutes until micturition or immediately if the pad became contaminated with faeces. Once the child had urinated, and wearing disposable gloves, the research nurse or clinical study officer (RN/CSO) removed the pad and urine was extracted into a sterile container as per the manufacturer's instructions. If it was not possible to obtain a sample prior to the child leaving the primary care site, the parent was given the necessary equipment and advice on obtaining a urine sample at home. The parent was advised to store the sample in the fridge and return it to their primary care site as soon as possible, ideally within 24 hours. The RN/CSO telephoned parents the next day to remind them to return the sample. Where feasible, the RN/CSO offered to collect the urine sample from the child's home.

At the primary care site, urine samples were dipstick tested (using Siemens/Bayer multistix 8SG) for blood, protein, glucose, ketones, nitrite, leukocyte esterase, pH and specific gravity (eight dipstick index tests). All index tests and the clinician's working diagnosis ('clinical diagnosis') were measured blind to the reference standard.

When there was at least 1ml of urine leftover urine after a priority sample was sent to a National Health Service (NHS) laboratory, it was sent using first class Royal Mail Safeboxes™ in boric acid monovettes to the Public Health Wales Microbiology Laboratory in Cardiff ("research laboratory"). Results from the research laboratory are used in the current analyses.

#### *Reference standard*

The research laboratory spiral plated (Don Whitley, UK) 50uL onto chromogenic agar and standard blood agar. A full description of the methods used in the research laboratory, and how these differed from the range of standard operating procedures in local NHS labs, is forthcoming.(10) Quantitative total counts were recorded for up to six organisms and the presence of antimicrobial substances measured. Samples were processed using a single, standardised procedure. We defined uropathogens as members of the *Enterobacteriaceae* group. Our microbiological definition of UTI was the presence of  $\geq 10^5$  Colony Forming Units (CFU)/mL of a single uropathogen ('pure growth') or  $\geq 10^5$  CFU/mL of a uropathogen with  $\geq 3 \log_{10}$

(1000-fold) difference between the growth of this and the next species ('predominant growth'). Aside from children's dates of birth, laboratory staff were blind to index tests. As there is no single accepted definition of contamination, we considered three definitions: growth of more than two organisms of more than  $>10^5$  (frank contamination) (11); growth of two or more organisms at more than  $10^5$  (12, 13); and growth of more than two organisms at greater than  $10^4$   $>2$  organisms (probable contamination according to Jackson(11) or frank contamination according to Bekeris(14)).

### *Statistical analysis*

We examined the frequency of symptom and sign categories, blind to their associations with urine culture results, and merged the least frequent categories prior to analyses. We used logistic regression to estimate associations of index tests with urine culture positivity. P-values were derived using likelihood ratio tests. For ordinal variables, both heterogeneity and trend p-values were derived. Continuous variables were divided into quintiles and trend p-values were derived using the median within categories. We examined plots of the log odds of culture positivity against the median within quintiles for evidence of non-linearity. We used two methods for dealing with missing data, including the response "don't know" to questions about the presence of symptoms such as pain, or crying when passing urine. In both univariable and multivariable analyses missing data were coded as the modal value, usually absence of the symptom. We repeated multivariable analyses using the chained equations approach to multiple imputation: estimates and Wald p-values (15) based on 50 imputed datasets derived using Rubin's rules.(16)

We derived prediction rules in three stages. First, we selected symptoms and signs with either trend or heterogeneity univariable p values  $<0.01$ . Second, we derived models from selected symptoms and signs using backwards stepwise selection and an exclusion criterion of heterogeneity p value  $>0.1$ . Third, we used backwards stepwise selection with the same exclusion p value for models in which dipstick results were added. We investigated the effect of using more liberal p value thresholds of 0.1 and 0.2 at the first stage, and found no important differences in the final models (results available from the authors on request).

We quantified diagnostic accuracy as the Area Under the Receiver Operating Characteristic (AUROC) curve (also known as the c-statistic). We also estimated AUROC values for clinical judgement of UTI. We conducted internal validation of the models using the bootstrap procedure described by Steyerberg (17) and calculated a calibration slope (shrinkage factor) by which we multiplied model coefficients in order to derive internally validated odds ratios. For each model, we selected cut-points corresponding to a range of values for sensitivity, and then calculated the corresponding specificity, negative and positive predictive values, and the proportion of children classified positive. These were compared against 'clinical diagnosis'



of UTI (where clinicians considered UTI to be 'fairly' or 'very' certain). We re-ran models leaving out predictive features that might lack face validity.

#### *Sample size calculation*

Our sample size calculation assumed a candidate predictor with 10% prevalence and UTI prevalence of 2%. With 80% power and a two-sided alpha of 5%, 3000 urine sample results were required to detect an odds ratio of 2.4 while 3100 results would give a 95% CI with width 10% for an algorithm with 80% sensitivity. We originally proposed to recruit 4000 children, combining analyses for children with both clean catch and nappy pad samples, anticipating we would recover urines from 77.5% for algorithm derivation, and a further 2000 children for external validation. However, we did not anticipate the need to stratify and report separate analyses by urine collection method. For the present analysis, we therefore decided to use all available nappy pad results to derive the model, and to conduct internal bootstrap validation instead of external validation.

## **Results**

### *Participants*

We recruited participants from 233 primary care sites (225 GP practices, four Walk-in Centres and four paediatric EDs) across England and Wales between April 2010 and April 2012. Of 10138 children screened eligible, 1276 (12.6%) declined participation, 1684 (16.6%) could not be recruited for other reasons and 15 (0.15%) withdrew.

Of the 234 primary care sites taking part, 198 sites (85%) completed and returned at least one screening log to the study centres. These show that 7350 children were screened but not recruited because they declined (1276), were not eligible (4390), or for other reasons (1684) which included: left the primary care site prior to invitation (811); did not give consent (214) or there was a language barrier (112) and an appropriate translator was not available at the time of recruitment. There were slightly more boys (mean difference of 5.2%; 95% CI 2.2% to 8.2%) among those for whom participation was declined ( $n = 1276$ ) compared to those who did agree to participate in the DUTY Study ( $n = 7163$ ). The mean age in the declined sample was 24.06 months vs. 26.88 months among participants (mean difference 2.04 months, 95% CI 1.08 to 3.00 months). We did not collect clinical information on those who declined.

Urine was collected from a total of 6241 children, 3205 using nappy pads. 3164 (98.7%) of nappy pad samples were sent to NHS labs and 2363 (73.7%) to the research laboratory. The number of reference standard results available from the research laboratory (our final analytic sample) was 2277 (71.0%). 82% of children providing nappy pad samples were aged <2 years (mean 1.3, SD 0.8), and the mean illness severity score was 2.3 points (SD 1.5). 48% of children were girls; and overall, 1.3% had a UTI (Table 1). The clean

### *Nappy pad urine sampling 03 February 2016*

catch sample (n=2740) had mean age 3.5 years (SD 1.0), mean illness severity score 2.2 points (SD 1.6) and 53.8% were girls.

2102 (92.3%) samples were provided within 24 hours of index test measurement and there was no relationship between UTI and time from urine collection to laboratory arrival. Antimicrobial substances, which can arise from the use of both systemic antibiotics and locally applied cleaning agents, were found in 6.6% of nappy pad samples, and were more likely to be present in children with than without UTI. GPs' 'working diagnosis' would have correctly identified 4 (13.3%) of the 30 UTIs with 97.0% specificity and an AUROC 0.63 (95% CI 0.53 to 0.72).

### *Nappy pad model*

Table 2 shows adjusted odds ratios for the index tests (marked\* in Table 1) selected for the nappy pad model. Parent-reported smelly urine, darker urine, female gender and the absence of a nappy rash were independently associated with UTI: for the first two there was evidence of graded associations. No clinical examination findings were independently associated with UTI. The presence of leukocytes and nitrites from dipstick urine testing were independently associated with UTI. The symptoms and signs model had reasonable diagnostic accuracy (validated AUROC for the multiple imputed model was 0.78 and diagnostic accuracy increased ( $p=0.036$ ) with addition of dipstick findings (validated AUROC 0.82)). Figure 2 shows the multiple imputed ROC curves for the models with and without dipstick urinalysis.

We re-ran the multiple imputation analysis excluding the nappy rash variable, and found that there was a reduction of 0.07 in validated AUROC in the symptom and sign model (from 0.78 to 0.71) and a reduction of 0.03 in the validated AUROC in the symptom, sign and dipstick model (from 0.82 to 0.80). We checked the association between antimicrobial substances in the urine and nappy rash, and found no association ( $p=0.82$ ).

### *Comparison of findings using nappy pads and catch samples*

The validated AUROC for the nappy pad model was inferior to the model derived using clean catch samples, which was 0.87 for symptoms and signs, increasing to 0.90 with dipstick results (to be reported fully in a future publication). Table 3 provides the proportion of nappy pad and clean catch samples considered to be contaminated according to three published definitions. (11-14) 'Frankly contaminated' urine was found in 12.2% of nappy pad and 1.8% of clean-catch samples, risk ratio 6.66 (95% CI 4.95 to 8.96;  $p < 0.001$ ).

## **Discussion**

*Summary of findings*

Four features (female gender, smelly urine, darker urine, and absence of nappy rash) that could be reported by parents, and no clinical signs, were associated with a microbiological diagnosis of UTI in children sampled using nappy pads. These features were substantially more predictive of a microbiological diagnosis of a UTI than clinicians' 'working' or clinical diagnoses, but less predictive than data obtained for older children sampled using the clean catch methods. More than 10% of samples obtained by nappy pads were 'frankly' contaminated, compared to less than 2% of samples obtained by clean catch from predominantly older children. The addition of dipstick testing improved diagnostic accuracy of nappy pad samples.

*Strengths and limitations*

The DUTY study is the largest primary care diagnostic accuracy study of clinical symptoms, signs and dipstick tests for diagnosing UTI in young children, and it achieved high levels of data completeness. We asked clinicians to obtain a clean catch urine sample whenever possible, but ultimately the decision whether to sample by clean catch or the nappy pad method was up to the parents, who generally used the nappy pad method in younger children. Children sampled using the nappy pad method were therefore younger and may have differed in other ways as well, for example being more unwell or distressed. We asked parents to replace pads at regular intervals until a sample was obtained. However, we did not use alarms to trigger scheduled replacement of pad, and this may have led to increased contamination rates.(12)

There were a relatively small number of UTI diagnoses, and we found that overall NHS laboratories have identified higher numbers of UTIs in this and in previous studies.(2, 18) It is plausible that the nappy pad contamination masked the presence of UTI leading to under-diagnosis in comparison to clean catch samples, in which lower contamination and higher UTI rates were observed. Our conservative criteria for a microbiological UTI diagnosis may have also contributed.

Our study reference standard defined uropathogens as members of the *Enterobacteriaceae* family at the UK guidelines'(19, 20) threshold of a pure/predominant growth of  $\geq 10^5$  CFU/ml. We chose not to use a lower threshold as this carries an increased risk of false positives, although there are recommendations that a lower threshold should be used.(3, 21, 22) The diagnosis of UTI is a clinical one taking microbiological analysis into account, and a lower microbiological threshold in the presence of high clinical suspicion would be acceptable for the purposes of clinical care as opposed to this diagnostic study. We used a rigorous criterion (minimum 3-log difference between the predominant and next most concentrated organism) for defining predominance. This definition could have reduced estimated prevalence if some UTIs were

incorrectly classified as contamination. In addition, a small proportion of the positive cultures may have been false positives due to asymptomatic bacteriuria or contamination.

Collecting an uncontaminated urine specimen is most difficult in the youngest children, and no method has yet been found to reliably distinguish pathogen from contaminant, especially when they co-exist. Our definition of UTI excluded atypical bacteria causing UTIs, which are also thought to be more common in younger children, potentially reducing our estimated UTI prevalence.(19) (23)

#### *Results in context with other studies*

The NICE guidelines found “insufficient data to draw conclusions about urine collection bags and urine collection pads,” but recommended their use when a clean catch sample cannot be obtained.(3) We have not been able to identify further studies addressing nappy pads since the NICE recommendations were published.(24) A systematic review of the accuracy of specimens obtained from nappy pads included three studies that compared sampling by nappy pad to sampling by urine bag, and one study that compared nappy pad specimens to specimens obtained by suprapubic aspiration. The latter study found 100% sensitivity and 94% specificity between the two methods. A randomised trial found that replacing pads every 30 minutes until a sample was obtained reduced contamination.(12)

We found a 1.3% prevalence of microbiological diagnoses of UTI. The only other UK primary care study found a 6% prevalence when urines were analyzed in National Health Service (NHS) laboratories.(2) We found a similar UTI prevalence of 5.6% for the DUTY study urines overall in NHS laboratories.(18) Fever was not an inclusion criterion in that study or in ours. However, a systematic review of 10 studies, eight of which were conducted in hospital emergency department, with one in a clinic and emergency department setting, and one in a clinic setting, and all conducted in the USA apart from a clinic study In Taiwan, found a UTI prevalence of 7% among infants presenting with fever,(25). An Australian study (with incomplete urine sampling) found a prevalence of 3.4% children presenting with fever to emergency departments. (26)

Our systematic search identified one systematic review that included eight primary studies of 7892 children aged less than 5 year(24) and three further primary studies (26-28) that included 17462 children, that assessed associations between clinical features symptoms and signs and a UTI diagnosis. The data we found show that no individual symptom or sign or combination of symptoms or signs was sufficient to rule in a diagnosis of UTI. Among the remaining studies, largely conducted in hospital emergency departments, abdominal pain, back pain, dysuria, frequency, and new-onset urinary incontinence increased the likelihood of a UTI.(29) Stridor, audible wheeze, circumcision, temperature <39°C with a source, abnormal chest sounds, chest crackles, age 3 years or less, not feeling hot, and breathing difficulty decreased the likelihood

of UTI. The largest study, which included almost 16,000 children aged less than five years presenting to emergency departments in Australia,(26) derived a diagnostic model based on a combination of 27 symptoms and signs. However, this study did not involve systematic urine sampling, and most children did not have urine sampled. This model was found to have an AUROC of 0.80 (95% CI 0.78 to 0.82), which is similar to findings from our DUTY study for children sampled with nappy pads.

Previous investigation of malodorous urine has shown conflicting results,(30) but our study strongly supports its diagnostic value. We investigated, but did not find evidence for, a number of non-specific symptoms (including fever, vomiting, lethargy, irritability and poor feeding) previously found to be associated with UTI (24) and recommended for clinical use by NICE.(3) It remains possible that such symptoms are of use in the secondary care settings in which studies reporting their utility were conducted, or in children with a different illness spectrum. This finding underlines the importance of including a wide range of illness presentation in studies of predictors of diagnoses, especially when symptoms and signs are notoriously non-specific. Studies that include only children with symptoms and signs previously found to be associated with the diagnosis risks missing previously unidentified predictors and 'research circularity' (looking for, and finding, symptoms and signs in children included in studies if they have those symptoms and signs).

The reduction in the risk of UTI associated with presence of a nappy rash should be interpreted with caution. The inverse association may arise through lower likelihood of a UTI when there is a plausible alternative diagnosis (conditioning on the common effect of primary care attendance). (31) Alternative explanations are that rash may be a risk factor for contamination of urine, and this masks the presence of a UTI, or that skin products used to treat nappy rash could render the urine sterile. However, we found no evidence of an association between nappy rash and contamination, nor the presence of antimicrobial substances in the urine and nappy rash. An increased likelihood of contamination of nappy pad samples might also explain the more modest associations of symptoms and dipstick test results with UTI than were found in clean catch samples.(10) We re-ran the models excluding nappy rash and found modest reductions in the symptoms and signs, and symptoms, signs and dipstick model AUROCs (respectively 0.07 and 0.03).

### *Conclusions*

Nappy pad urine sample culture results together with symptoms that can be reported by parents can be clinically useful in identifying acutely unwell pre-school children presenting to primary care who have a UTI, but with less accuracy, and with increased contamination compared to clean catch sampling. Clean catch urine sampling in children in primary care should be prioritised, especially in children with nappy rash.

However, if sampling is done using nappy pads, then the addition of dipstick testing significantly improves diagnostic accuracy.

Further research is needed to distinguish pathogens from contaminants, and to establish the cost effectiveness of different sensitivity/specificity cut points using routine health service laboratory results. We do not know precisely how results from clean catch sampling would compare to nappy pad sampling in younger children: contamination may vary by age as well as by sampling method. Randomising children to sampling method would shed light on this.

*Ethical approval and the role of the funder*

Multi-centre ethical approval was granted by the South West Southmead Research Ethics Committee (previously Southmead Research Ethics Committee, then South West 4 REC), Ref #09/H0102/64.

This project was funded by the National Institute for Health Research (NIHR) Health Technology Assessment (HTA) programme (project number 08/66/01) and will be published in full in Health Technology Assessment. Further information available at: [www.DUTYstudy.org.uk](http://www.DUTYstudy.org.uk)

This report presents independent research commissioned by the National Institute for Health Research (NIHR). The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, MRC, CCF, NETSCC, the National Institute for Health Research (NIHR) Health Technology Assessment (HTA) programme or the Department of Health.

The corresponding author confirms he had full access to all data and takes final responsibility for submitting the manuscript for publication.

*Competing interest statement*

All authors declare they have no conflict of interest and have signed the conflict of interest form.

*Acknowledgements*

We wish to thank the children, their families, the GP practices and other recruitment sites, the research networks, the NHS laboratories and all the members of the DUTY team including: Steven Beech, Jonathan Benger, Theresa Bowes, Peter Brindle, Lisa Patten, Christina Curry, Lewis Darmanin, Catherine Derrick, Micaela Gal, Susan George, Margaret Hague, Andrea Jarman, Lyn Liddiard, Ruth Munn, Marilyn Peters, Carolyn Powell, Jennifer Richards, Victoria Roberts, Annie Sadoo, Elizabeth Thomas, Tessa Wade, Stana Williams and Jane Woodhead. We also wish to thank the providers of nursing/clinical studies officer support from the Primary Care Research Networks in: Greater London; Kent and Medway; Sussex; Surrey; Thames Valley; the South-west; Cumbria and Lancashire; Northumberland and Tyne and Wear; and the National Institute for Social Care and Health Research, Clinical Research Centre in Wales (NISCHR-CRC). Additionally, we wish to acknowledge the support given by the South East Wales Trials Unit (funded by NISCHR), Comprehensive Local Research Networks of Central and East London, Western, Peninsula, Hampshire and Isle of Wight, and the NIHR Biomedical Research and Development Department, Guy's and St Thomas' NHS Foundation Trust.

*Nappy pad urine sampling 03 February 2016*

We thank Study Steering Committee members: Frank Sullivan, Rafael Perera, Matthew Thompson and Clodna McNulty.



Table 1. Nappy pad samples: children's characteristics and crude odds ratios for index tests\* associated with UTI

Demographics/index tests	N (%) <sup>a</sup>	UTI (%) <sup>b</sup>	Crude OR (95% CI) <sup>c</sup>
<b>Total</b>	2277	30 (1.3%)	
<b>Age of child</b>			
< 6 months	369 (16.2%)	5 (1.4%)	1.72 (0.54,5.46)
6 - < 12 months	603 (26.5%)	11 (1.8%)	2.33 (0.90,6.04)
1 - < 2 years	884 (38.8%)	7 (0.8%)	1 (ref)
2 - <3 years	353 (15.5%)	7 (2.0%)	2.53 (0.88, 7.28)
3 - <4 years	58 (2.5%)	0 (0.0%)	n/a
4 years plus	10 (0.4%)	0 (0.0%)	n/a
<b>Time from index tests to taking urine sample</b>			
Sample before recruitment	120 (5.3%)	2 (1.7%)	1.33 (0.31,5.67)
< 24 hours	1982 (87.0%)	25 (1.3%)	1 (ref)
24 hours to < 48 hours	109 (4.8%)	3 (2.8%)	2.22 (0.66,7.45)
48 hours to < 72 hours	18 (0.8%)	0 (0.0%)	n/a
72 hours plus	48 (2.1%)	0 (0.0%)	n/a
<b>Clinician diagnosis prior to dipstick</b>			
No UTI certain to very certain	1033 (45.4%)	8 (0.8%)	0.52 (0.22,1.19)
Uncertain or No UTI fairly certain	1201 (52.7%)	18 (1.5%)	1(ref)
UTI fairly to very certain	38 (1.7%)	4 (10.5%)	7.76 (2.49, 24.18)
Missing	5 (0.2%)	0 (0.0%)	
<b>Gender</b>			
Male	1183 (52.0%)	9 (0.8%)	1 (ref)
Female	1094 (48.0%)	21 (1.9%)	2.55 (1.16,5.60)
<b>Smelly urine</b>			
No problem	1518 (66.7%)	12 (0.8%)	1 (ref)
Slight problem	206 (9.0%)	4 (1.9%)	2.20 (0.73,6.61)
Moderate problem	138 (6.1%)	5 (3.6%)	4.18 (1.52,11.50)
Severe problem	26 (1.1%)	4 (15.4%)	20.21 (6.29,64.97)
Missing/not known	389 (17.1%)	5 (1.3%)	
<b>Darker urine</b>			
No problem	1764 (77.5%)	19 (1.1%)	1 (ref)
Slight problem	83 (3.6%)	2 (2.4%)	2.19 (0.51,9.43)
Moderate or severe problem	41 (1.8%)	4 (9.8%)	9.59 (3.17,29.02)

*Nappy pad urine sampling 03 February 2016*

Demographics/index tests	N (%) <sup>a</sup>	UTI (%) <sup>b</sup>	Crude OR (95% CI) <sup>c</sup>
Missing/not known	389 (17.1%)	5 (1.3%)	
<b>Nappy rash<sup>d</sup></b>			
No problem	1715 (75.3%)	29 (1.7%)	1 (ref)
Slight to severe problem	560 (24.6%)	1 (0.2%)	0.10 (0.01,0.77)
Missing	2 (0.1%)	0 (0.0%)	
Dipstick: leucocytes			
Negative	1759 (77.3%)	13 (0.7%)	1 (ref)
Trace	125 (5.5%)	1 (0.8%)	1.09 (0.14,8.38)
+	119 (5.2%)	4 (3.4%)	4.69 (1.50,14.61)
++	177 (7.8%)	4 (2.3%)	3.12 (1.01,9.66)
+++	91 (4.0%)	8 (8.8%)	12.99 (5.24,32.20)
Missing	6 (0.3%)	0 (0.0%)	
Dipstick: nitrites			
Negative	1916 (84.1%)	13 (0.7%)	1 (ref)
Positive	355 (15.6%)	17 (4.8%)	7.39 (3.55,15.35)
Missing	6 (0.3%)	0 (0.0%)	

<sup>a</sup> Percentage relative to the total number of observations (i.e. 2277)

<sup>b</sup> Percentage relative to the total number of observations within that category

<sup>c</sup> Crude ORs calculated using modal imputation

Table 2. Nappy pad samples: models based on symptoms and signs; and on symptoms, signs and dipstick results, including results based on multiple imputation. Odds ratios calculated using shrunken estimates from the bootstrap internal validation calibration slope

Index tests	Symptom and sign model		Symptom sign and dipstick model	
	Adjusted OR (95 % CI) <sup>a</sup>	MI <sup>b</sup> adjusted OR (95% CI)	Adjusted OR (95 % CI) <sup>a</sup>	MI <sup>b</sup> adjusted OR (95% CI)
<b>Gender</b>				
Male	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Female	1.95 (1.11,3.41)	1.96 (1.06,3.61)	1.41 (0.80,2.48)	1.45 (0.78,2.72)
<b>Smelly urine</b>				
No problem	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Slight problem	1.61 (0.73,3.54)	1.97 (0.82,4.71)	1.44 (0.67,3.11)	1.79 (0.76,4.23)
Moderate problem	2.51 (1.14,5.51)	3.39 (1.46,7.89)	2.15 (0.98,4.68)	2.96 (1.26,6.97)
Severe problem	7.40 (2.98,18.36)	10.14 (3.85,26.69)	3.97 (1.58,9.96)	6.13 (2.28,16.47)
<b>Darker urine</b>				
No problem	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Slight problem	1.89 (0.66,5.46)	1.99 (0.65,6.12)	1.81 (0.65,5.07)	1.92 (0.63,5.88)
Moderate or severe problem	2.46 (0.98,6.21)	2.26 (0.85,6.01)	2.29 (0.93,5.62)	2.27 (0.87,5.93)
<b>Nappy rash</b>				
No problem	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Slight to severe problem	0.16 (0.04,0.66)	0.13 (0.03,0.61)	0.19 (0.05,0.71)	0.16 (0.04,0.66)
<b>Dipstick: leukocytes</b>				
Negative			1 (ref)	1 (ref)
Trace			0.87 (0.23,3.31)	0.81 (0.18,3.61)
+			2.06 (0.92,4.63)	2.18 (0.88,5.43)
++			1.63 (0.73,3.62)	1.78 (0.73,4.30)
+++			3.27 (1.66,6.41)	3.35 (1.57,7.15)
<b>Dipstick: nitrites</b>				
Negative			1 (ref)	1 (ref)
Positive			3.16 (1.91,5.24)	3.70 (2.10,6.52)
<b>ROC (95% CI)</b>	0.769 (0.68, 0.85)	0.805 (0.72, 0.89)	0.858 (0.79, 0.93)	0.870 (0.80, 0.94)
<b>Validated ROC<sup>c</sup></b>	0.744	0.778	0.799	0.821
<b>Δ ROC<sup>d</sup> (95% CI)</b>			0.089 (0.02,0.16)	0.065 (0.00,0.13)
<b>Δ ROC<sup>d</sup> p-value</b>			0.012	0.036

Calibration slope	0.695	0.749	0.647	0.708
-------------------	-------	-------	-------	-------

<sup>a</sup> Missing values coded to modal category <sup>b</sup>MI: multiple imputation <sup>c</sup> Internal validation using the bootstrap procedure <sup>d</sup> The difference in ROC between symptom and sign model and symptom, sign and dipstick model

Figure 2. Nappy pad ROC curve from multiple imputation symptoms and signs only (solid line) and symptoms, signs and dipstick (dotted line)




Table 3. The numbers of contaminated samples using different definitions of contamination. The highlighted row is the definition of contamination that we use in our subsequent analyses

Paper	Contamination definition	Clean catch contaminated N(%)	Nappy pad contaminated N(%)	Risk difference (95% CI)	Risk ratio (95% CI)	P value
Jackson (11), frank contamination	>10e5 >2 organisms	50/2740 (1.8%)	277/2277 (12.2%)	0.103 (0.089,0.118)	6.666 (4.959,8.963)	<0.001
Feasey (13)or Rao(12)	>10e5 ≥2 organisms	78/2740 (2.8%)	426/2277 (18.7%)	0.159 (0.141,0.176)	6.572 (5.196,8.312)	<0.001
Jackson(11) probable or frank contamination or Bekeris(14)	>10e4 >2 organisms	175/2740 (6.4%)	599/2277 (26.3%)	0.199 (0.179,0.219)	4.119 (3.513,4.829)	<0.001

## Reference

1. Coulthard MG, Vernon SJ, Lambert HJ, et al. A nurse led education and direct access service for the management of urinary tract infections in children: prospective controlled trial. *BMJ*. 2003;327(7416):656.
2. O'Brien K, Edwards A, Hood K, et al. Prevalence of urinary tract infection in acutely unwell children in general practice: a prospective study with systematic urine sampling. *Br J Gen Pract*. 2013;63(607):e156-64.
3. National Institute for Health and Care Excellence (NICE). Urinary Tract Infection in Children: diagnosis, treatment and long-term management. NICE Clinical Guideline 54. 2007.
4. van der Voort J, Edwards A, Roberts R, et al. The struggle to diagnose UTI in children under two in primary care. *Fam Pract*. 1997;14(1):44-8.
5. Liaw LC, Nayar DM, Pedler SJ, et al. Home collection of urine for culture from infants by three methods: survey of parents' preferences and bacterial contamination rates. *BMJ*. 2000;320(7245):1312-3.
6. Vernon S, Foo CK, Coulthard MG. How general practitioners manage children with urinary tract infection: an audit in the former Northern Region. *Br J Gen Pract*. 1997;47(418):297-300.
7. Adams ST, Leveson SH. Clinical prediction rules. *BMJ*. 2012;344:d8312.
8. Downing H, Thomas-Jones E, Gal M, et al. The diagnosis of urinary tract infections in young children (DUTY): protocol for a diagnostic and prospective observational study to derive and validate a clinical algorithm for the diagnosis of UTI in children presenting to primary care with an acute illness. *BMC Infect Dis*. 2012;12:158.
9. Vernon S, Redfearn A, Pedler SJ, et al. Urine collection on sanitary towels. *Lancet*. 1994;344(8922):612.
10. Hay AD, Birnie K, Busby J, et al. The Diagnosis of Urinary Tract Infections in Young Children (DUTY): a diagnostic and prospective observational study to derive and validate a clinical algorithm for the diagnosis of UTI in children presenting to primary care with an acute illness. *Health Technol Assess*. In press.
11. Jackson SR, Dryden M, Gillett P, et al. A novel midstream urine-collection device reduces contamination rates in urine cultures amongst women. *BJU Int*. 2005;96(3):360-4.
12. Rao S, Bhatt J, Houghton C, et al. An improved urine collection pad method: a randomised clinical trial. *Archives of disease in childhood*. 2004;89(8):773-5.
13. Feasey S. Are Newcastle urine collection pads suitable as a means of collecting specimens from infants? *Paediatr Nurs*. 1999;11(9):17-21.
14. Bekeris LG, Jones BA, Walsh MK, et al. Urine culture contamination: a College of American Pathologists Q-Probes study of 127 laboratories. *Arch Pathol Lab Med*. 2008;132(6):913-7.
15. Kinlin LM, Kirchner C, Zhang H, et al. Derivation and validation of a clinical prediction rule for nosocomial pneumonia after coronary artery bypass graft surgery. *Clin Infect Dis*. 2010;50(4):493-501.
16. Rubin D. Inference and missing data. *Biometrika*. 1976;63(3):581-92.
17. Steyerberg EW, Harrell FE, Jr., Borsboom GJ, et al. Internal validation of predictive models: efficiency of some procedures for logistic regression analysis. *J Clin Epidemiol*. 2001;54(8):774-81.
18. Butler CC, O'Brien K, Pickles T, et al. Childhood urinary tract infection in primary care: a prospective observational study of prevalence, diagnosis, treatment, and recovery. *Br J Gen Pract*. 2015;65(633):e217-23.
19. Friedman S, Reif S, Assia A, et al. Clinical and laboratory characteristics of non-E. coli urinary tract infections. *Arch Dis Child*. 2006;91(10):845-6.
20. HealthProtectionAgency. National standard method: investigation of urine (BSOP 41). 2009 2009. Report No.
21. Grabe M, Bjerklund-Johansen TE, Botto H, et al. Guidelines on Urological Infections: European Association of Urology; 2013. Available from: [http://uroweb.org/wp-content/uploads/18\\_Urological-infections\\_LR.pdf](http://uroweb.org/wp-content/uploads/18_Urological-infections_LR.pdf).
22. Scottish Intercollegiate Guidelines Network (SIGN). Management of suspected bacterial urinary tract infection in adults (SIGN publication no.88) Edinburgh: SIGN; 2012. Available from: <http://www.sign.ac.uk/pdf/sign88.pdf>.

23. Paschke AA, Zaoutis T, Conway PH, et al. Previous antimicrobial exposure is associated with drug-resistant urinary tract infections in children. *Pediatrics*. 2010;125(4):664-72.
24. Hay AD, Whiting P, Butler CC. How best to diagnose urinary tract infection in preschool children in primary care? *BMJ*. 2011;343:d6316.
25. Shaikh N, Morone NE, Bost JE, et al. Prevalence of urinary tract infection in childhood: a meta-analysis. *Pediatr Infect Dis J*. 2008;27(4):302-8.
26. Craig JC, Williams GJ, Jones M, et al. The accuracy of clinical symptoms and signs for the diagnosis of serious bacterial infection in young febrile children: prospective cohort study of 15 781 febrile illnesses. *BMJ*. 2010;340:c1594.
27. Gorelick MH, Shaw KN. Clinical decision rule to identify febrile young girls at risk for urinary tract infection. *Arch Pediatr Adolesc Med*. 2000;154(4):386-90.
28. Armengol CE, Hendley JO, Schlager TA. Should we abandon standard microscopy when screening for urinary tract infections in young children? *Pediatr Infect Dis J*. 2001;20(12):1176-7.
29. Shaikh N, Morone NE, Lopez J, et al. Does This Child Have a Urinary Tract Infection? *JAMA: The Journal of the American Medical Association*. 2007;298(24):2895-904.
30. Shaw KN, Hexter D, McGowan KL, et al. Clinical evaluation of a rapid screening test for urinary tract infections in children. *J Pediatr*. 1991;118(5):733-6.
31. Lejeune B, Baron R, Guillois B, et al. Evaluation of a screening test for detecting urinary tract infection in newborns and infants. *J Clin Pathol*. 1991;44(12):1029-30.